

Notes on the water bugs (Hemiptera: Heteroptera) in urban areas of Hanoi

Tran Anh Duc*, Pham Thi Mai, Nguyen Xuan Quynh,
Ngo Xuan Nam, Nguyen Quang Huy

Faculty of Biology, Hanoi University of Science, VNU, 334 Nguyen Trai, Hanoi, Vietnam

Received 28 December 2010

Abstract. In the present study, faunistic data on water bugs from urban area of Hanoi has been catalogued for the first time. From our surveys, we have identified a total of 23 species of 12 genera and nine families of aquatic bugs (Nepomorpha) and semi-aquatic bugs (Gerromorpha) from the studied area. The water bug fauna in Hanoi is of a typical lowland fauna in Southeast Asia mainland, comprising many species with wide distribution. Several taxa are expected to exist in the studied area, but they have not been seen during our field collection, suggesting further field studies are needed.

Keywords: water bugs, Hemiptera, Gerromorpha, Nepomorpha, Hanoi.

1. Introduction

Hanoi is situated in the centre of the Red River delta, in the North of Vietnam, with a typical humid subtropical climate. Hanoi is a city between rivers, thus has many natural lakes (which are traces of ancient rivers). Its central town has several “rivers” which are in fact channels created few hundreds years ago, such as To Lich River and Nhue River. Natural rivers in Hanoi run mostly through its rural districts [1]. Recent developments have caused various levels of pollution in Hanoi area, especially in the central town [1]. This certainly affects the lives of aquatic organisms like water bugs (order Hemiptera).

The water Hemiptera comprises two infra-orders, namely Gerromorpha (semi-aquatic bugs) and Nepomorpha (true aquatic bugs). The fauna of Vietnamese water bugs has been studied over past few decades, mainly by foreign scientists. Those studies mostly based on samples collected from natural habitats. To date, there has yet any study on aquatic and semi-aquatic bugs in urban water bodies. There have been only few incidental records of aquatic bugs from Hanoi in published literature, i.e., *Hydrometra albolineata*, *H. annamana* [2].

This study aims to catalogue the water bug fauna in the central town of Hanoi. The results from this study are to be served as the first base-line data for subsequent studies on the fauna and ecology of water bugs from the urban area.

* Corresponding author. Tel.: 84-4-38582795.
E-mail: tran.anhduc@hus.edu.vn

2. Materials and methods

Field surveys were conducted during the period of April - November 2010 from various localities in the central town of Hanoi. Samples were collected by hand nets and pond nets from selected water bodies, representing still waters (ponds, lakes, and paddy-fields) and running ones (artificial rivers). The following list of sampling localities contains the collecting codes used also in the results and discussion section.

(a) Still waters:

- Ponds, lakes:

HN01: Lotus pond 1 near West Lake Water Park (left)

HN02: Lotus pond 2 near West Lake Water Park (right)

HN03: West Lake

HN04: Thanh Huong Lake (used for recreational fishing), near Khuat Duy Tien road, (Thanh Xuan district)

HN05: Linh Dam Lake

HN06: Anh Duong Lake (used for recreational fishing), Me Tri area (Thanh Xuan district)

- Paddy-fields:

HN11: Paddy-field at Thuan village, Phu Lam ward (Ha Dong district)

HN13: Paddy-field at Dai Mo area (Tu Liem district)

HN14: Paddy-field at Ngoc Truc area (Tu Liem district)

HN15: Paddy-field near Noi Bridge of Nhue River

(b) Running waters:

HN07: Nhue River, at Doi Bridge

HN08: Nhue River, at Lien Mac area

HN09: Nhue River, at Noi Bridge

HN10: To Lich River

HN12: Day River, at Mai Linh Bridge (Ha Dong district)

Voucher specimens are deposited in the Hydrobiology Laboratory, Department of Invertebrate Zoology, Hanoi University of Science. Specimens collected were kept in 70 % ethanol.

Identification of various water bug groups were mainly based on the keys provided by Chen *et al.* (2005), and other studies on each respective group, i.e. Lansbury (1972) for *Ranatra* (Nepidae); Nieser (2002) for Micronectidae; Nieser (2004) for Notonectidae and Pleidae; Andersen *et al.* (2002) for *Microvelia* (Veliidae); Andersen (1990), Cheng *et al.* (2001) for Gerridae [3-11].

3. Results and discussion

From our surveys, we have identified a total of 23 species of 12 genera and nine families of aquatic bugs (infra-order Nepomorpha) and semi-aquatic bugs (infra-order Gerromorpha) from the studied area (Table 1). Among these species, six species could only be keyed down to generic level, because these are poorly studied genera and species identities are difficult to confirm. *Laccotrephes* specimen is a nymph, thus not possible to be identified to species level. Most species found in the studied area have wide distribution in Southeast Asia, and commonly found at lowland area [3-11]. The most common species at various habitats are *Aquarius padulum padulum*, *Mesovelia vittigera*, *Microvelia douglasi*, *Diplonychus rusticus*, *Micronecta quadristrigata*, *Micronecta ludibunda*, and *Anisops kuroiwae*.

Species diversity in different habitats: We have observed better species richness and higher abundance of aquatic bugs in certain sampling sites, i.e. site number HN01, HN02, HN04, HN06, HN12. These sites are relatively

“natural and clean”, meaning to have little artificial impact and relatively good aquatic vegetation (or submerged grass). In other sites with significantly lower species richness and abundance, we have observed certain levels of artificial disturbance, e.g. pollution, agricultural work, concretised banks, and absence (or removal) of aquatic vegetation. From our observations, we believe that the presence of aquatic vegetation significantly contribute to the richness of water bug fauna. Several studies have already reported that aquatic vegetation provides shelters and resting places for many water bug species [3].

We could not find any water bug sample from To Lich River (HN10) probably due to heavy pollution there. This river is the main drainage of wastewater for the city. The pollution here is probably over the limit that aquatic and semi-aquatic bugs could tolerate.

Missing taxa: The first author has previously examined the collection of the Budapest Natural History Museum and found specimens collected from the central town of Hanoi. They belong to the following species *Hydrometra albolineata*, *H. annamana* [2], *Rhagadotarsus kraepelini*, *Neogerris parvulus*. These samples were collected by Hungarian scientists during 1960s-1980s, when Hanoi was much less urbanised than today. However, in recent surveys, we did not encounter any individual of these species. These species may vanish due to urbanisation and water pollution. However, they may still live around the rural districts of Hanoi, which is out of the scope of this study. Future field collection, both in the rural and urban parts, may help to re-confirm the occurrence of these species in Hanoi area.

Table 1. Aquatic Hemiptera collected from Hanoi city during field survey in 2010
(Notes: x = present; Site number HN10 is excluded as no water bug was found there)

No.	Taxon	Sampling localities	HN 01	HN 02	HN 03	HN 04	HN 05	HN 06	HN 07	HN 08	HN 09	HN 11	HN 12	HN 13	HN 14	HN 15
Infra-order Gerromorpha																
Family Gerridae																
1	<i>Aquarius paludum paludum</i> (Fabricius, 1794)		x	x		x		x		x		x	x			
2	<i>Limnomogus fossarum fossarum</i> (Fabricius, 1775)		x	x		x		x								
3	<i>Limnogonus nitidus</i> (Mayr, 1865)			x												
Family Hebridae																
4	<i>Hebrus</i> sp.			x										x		
Family Mesoveliidae																
5	<i>Mesovelia horvathi</i> Lundblad, 1933		x	x												
6	<i>Mesovelia vittigera</i> Horváth, 1915		x		x	x	x	x	x		x		x			
Family Veliidae																
7	<i>Microvelia douglasi</i> Scott, 1847		x	x			x						x	x	x	x
8	<i>Microvelia leveillei</i> (Lethierry, 1877)														x	
9	<i>Microvelia</i> sp.1					x	x						x	x		x
10	<i>Microvelia</i> sp.2			x		x	x						x	x		x
11	<i>Microvelia</i> sp.3			x		x	x						x		x	

No.	Taxon	Sampling localities														
		HN 01	HN 02	HN 03	HN 04	HN 05	HN 06	HN 07	HN 08	HN 09	HN 11	HN 12	HN 13	HN 14	HN 15	
Infra-order Nepomorpha																
Family Belostomatidae																
12	<i>Diplonychus rusticus</i> (Fabricius, 1871)	x	x		x	x	x					x			x	
Family Micronectidae																
13	<i>Micronecta quadristrigata</i> Breddin, 1905	x	x	x	x	x	x		x	x	x	x	x	x	x	x
14	<i>Micronecta ludibunda</i> Breddin, 1905	x	x	x	x		x		x	x		x				
15	<i>Micronecta scutellaris</i> (Stål, 1868)	x					x			x						
16	<i>Micronecta tarsalis</i> Chen, 1960								x							
Family Nepidae																
17	<i>Ranatra longipes longipes</i> Stål, 1861				x											
18	<i>Laccotrephes</i> sp.					x										
Family Notonectidae																
19	<i>Anisops breddini</i> Kirkaldy, 1901	x	x				x									
20	<i>Anisops kuroiwa</i> e Matsumura, 1905	x					x		x			x		x	x	
21	<i>Anisops</i> sp.				x		x					x		x	x	
22	<i>Nychia sappho</i> Stål, 1859	x	x				x									
Family Pleidae																
23	<i>Paraplea</i> sp.	x	x													

4. Conclusion

Based on our surveys, the fauna of water bugs in Hanoi is a typical lowland fauna in Southeast Asia mainland, comprising many species with wide distribution. Several taxa are expected to exist in the studied area, but they have not been seen during our field collection. Thus, further field studies are needed for more comprehensive understandings of water bug diversity in the studied area.

Acknowledgements

This study is supported by Research Grant number TN-10-29 of Hanoi University of Science (Vietnam National University).

References

- [1] An overview of Hanoi: <http://www.vietnamtravelco.com/dulich/kham-pha-du-lich-ha-noi.htm> (accessed 29 Mar. 2011)
- [2] A.D. Tran, C.M. Yang, X.Q. Nguyen & H. Zettel, Faunistical notes on the water measurer *Hydrometra* Latreille, 1796 (Insecta: Heteroptera: Hydrometridae) from Vietnam and Hainan Island, *Annalen des Naturhistorischen Museums in Wien* 111B (2010) 19.
- [3] P.P. Chen, N. Nieser, H. Zettel, *The aquatic and semi-aquatic bugs (Heteroptera: Nepomorpha & Gerromorpha) of Malesia*. Fauna Malesiana Handbooks 5, Brill, Leiden - Boston, 2005.
- [4] I. Lansbury, A review of the Oriental species of *Ranatra* Fabricius (Hemiptera-Heteroptera: Nepidae), *Transactions of the Royal Entomological Society of London* 124 (1972) 287.

- [5] N.M. Andersen, C.M. Yang & H. Zettel, Notes on the Microveliinae of Singapore and Peninsula Malaysia with the description of two new species of *Microvelia* Westwood (Hemiptera-Heteroptera: Veliidae), *Raffles Bulletin of Zoology* 50(1) (2002) 111.
- [6] N.M. Andersen, C.M. Yang & H. Zettel, Guide to the aquatic Heteroptera of Singapore and Peninsular Malaysia. 2. Veliidae, *Raffles Bulletin of Zoology* 50(1) (2002) 231.
- [7] L. Cheng, C.M. Yang & J.T. Polhemus, Guide to aquatic Heteroptera of Singapore and Peninsular Malaysia. I. Gerridae and Hermatobatidae, *Raffles Bulletin of Zoology* 49(1) (2001) 129.
- [8] N. Nieser, Guide to aquatic Heteroptera of Singapore and Peninsular Malaysia. IV. Corixoidea, *Raffles Bulletin of Zoology* 50(1) (2002) 263.
- [9] N. Nieser, Guide to aquatic Heteroptera of Singapore and Malay Peninsular Malaysia III. Pleidae and Notonectidae, *Raffles Bulletin of Zoology* 52(1) (2004) 79.
- [10] N.M. Andersen, Phylogeny and taxonomy of the water striders, genus *Aquarius* Schellenberg (Insecta, Hemiptera, Gerridae), with a new species from Australia, *Steenstrupia* 16(4) (1990) 37.
- [11] N.M. Andersen, The *Limnogonus* and *Neogerris* of the Old World with character analysis and a reclassification of the Gerrinae, *Entomologica Scandinavica* Suppl.7 (1975) 7.

Dẫn liệu về thành phần côn trùng nước thuộc bộ Hemiptera tại khu vực nội thành Hà Nội

Trần Anh Đức, Phạm Thị Mai, Nguyễn Xuân Quýnh,
Ngô Xuân Nam, Nguyễn Quang Huy

Khoa Sinh học, Trường Đại học Khoa học Tự nhiên, ĐHQGHN, 334 Nguyễn Trãi, Hà Nội, Việt Nam

Kết quả phân tích vật mẫu thu được trong các đợt khảo sát năm 2010 tại khu vực nội thành Hà Nội đã xác định được 23 loài thuộc 12 giống, 9 họ, 2 phân bộ (Gerromorpha và Nepomorpha) của Côn trùng nước bộ Hemiptera. Phần lớn những loài thu được tại khu vực nghiên cứu đều thuộc khu hệ vùng đồng bằng điển hình của Đông Nam Á, với nhiều loài có phân bố rộng. Tuy nhiên, cần có thêm những nghiên cứu đầy đủ hơn để có thể đánh giá chính xác hiện trạng côn trùng nước bộ Hemiptera tại khu vực Hà Nội cả ở khu vực nội thành và các huyện ngoại thành.