

The nonlinear absorption coefficient of a strong electromagnetic wave by confined electrons in quantum wells under the influences of confined phonons

Bau N.Q., Hung L.T., Nam N.D.

Department of Physics, College of Natural Sciences, Hanoi National University, No. 334, Nguyen Trai Str.,
Thanh Xuan Dist., Hanoi, Viet Nam

Abstract: The nonlinear absorption coefficient (NAC) of a strong electromagnetic wave (EMW) by confined electrons in quantum wells under the influences of confined phonons is theoretically studied by using the quantum transport equation for electrons. In comparison with the case of unconfined phonons, the dependence of the NAC on the energy ($\hbar\omega$), the amplitude (E_0) of external strong EMW, the width of quantum wells (L) and the temperature (T) of the system in both cases of confined and unconfined phonons is obtained. Two limited cases for the absorption: close to the absorption threshold ($|\hbar\omega - \hbar\omega_0| \leq \epsilon$) and far away from the absorption threshold ($|\hbar\omega - \hbar\omega_0| \gg \epsilon$) (ω_0 and ϵ are the frequency of optical phonon and the average energy of electron, respectively) are considered. The formula of the NAC contains the quantum number m characterizing confined phonons and is easy to come back to the case of unconfined phonons and linear absorption. The analytic expressions are numerically evaluated, plotted and discussed for a specific case of the GaAs/GaAsAl quantum well. Results show that there are more resonant peaks of the NAC which appear in the case of confined phonons when $\omega > \omega_0$ than in that of unconfined phonons. The spectrums of the NAC are very different from the linear absorption and strongly depend on m . © 2010 VSP.

Index Keywords: Analytic expressions; Average energy; Confined phonons; Linear absorption; Nonlinear absorption coefficient; Optical phonons; Quantum numbers; Quantum transport equations; Quantum well; Resonant peaks; Absorption; Cements; Electromagnetic wave diffraction; Electromagnetic wave scattering; Electromagnetic waves; Electromagnetism; Electrons; Nonlinear equations; Quantum chemistry; Quantum electronics; Semiconductor quantum wells; Phonons

Year: 2010

Source title: Journal of Electromagnetic Waves and Applications

Volume: 24

Issue: 13

Page : 1751-1761

Link: Scopus Link

Correspondence Address: Bau, N. Q.; Department of Physics, College of Natural Sciences, Hanoi National University, No. 334, Nguyen Trai Str., Thanh Xuan Dist., Hanoi, Viet Nam; email: nguyenquangbau54@gmail.com

ISSN: 9205071

CODEN: JEWAE

Language of Original Document: English

Abbreviated Source Title: Journal of Electromagnetic Waves and Applications

Document Type: Article

Source: Scopus

Authors with affiliations:

1. Bau, N.Q., Department of Physics, College of Natural Sciences, Hanoi National University, No. 334, Nguyen Trai Str., Thanh Xuan Dist., Hanoi, Viet Nam
2. Hung, L.T., Department of Physics, College of Natural Sciences, Hanoi National University, No. 334, Nguyen Trai Str., Thanh Xuan Dist., Hanoi, Viet Nam
3. Nam, N.D., Department of Physics, College of Natural Sciences, Hanoi National University, No. 334, Nguyen Trai Str., Thanh Xuan Dist., Hanoi, Viet Nam

References:

1. Rucker, H., Molinari, E., Lugli, P., Microscopic calculation of the electron-phonon interaction in quantum wells (1992) *Phys. Rev. B*, 45, p. 6747
2. Vasilopoulos, P., Charbonneau, M., Van Vliet, C.M., Linear and nonlinear electrical conduction in quasi-two-dimensional quantum wells (1987) *Phys. Rev. B*, 35, p. 1334
3. Suzuki, A., Theory of hot-electron magneto phonon resonance in quasi-two-dimensional quantum-well structures (1992) *Phys. Rev. B*, 45, p. 6731
4. Bau, N.Q., Phong, T.C., Calculations of the absorption coefficient of a weak electromagnetic wave by free carriers in quantum wells by the Kubo-Mori method (1998) *J. Phys. Soc. Jpn.*, 67, p. 3875
5. Bau, N.Q., Nhan, N.V., Phong, T.C., Calculations of the absorption coefficient of a weak electromagnetic wave by free carriers in doped superlattices by using the Kubo-Mori method (2002) *J. Korean. Phys. Soc.*, 41, p. 149
6. Schmittink, S., Chemla, D.S., Miller, D.A.B., Linear and nonlinear optical properties of semiconductor quantum wells (1989) *Adv. Phys.*, 38, p. 89
7. Ploog, K., Dohler, G.H., Compositional and doping superlattices in II-V semiconductors (1983) *Adv. Phys.*, 32, p. 285
8. Richter, M., Carmele, A., Butscher, S., Bücking, N., Milde, F., Kratzer, P., Scheffler, M., Knorr, A., Two-dimensional electron gases: Theory of ultrafast dynamics of electron-phonon interactions in graphene, surfaces, and quantum wells (2009) *J. Appl. Phys.*, 105, p. 122409
9. Shih, T., Reimann, K., Woerner, M., Elsaesser, T., Waldmüller, I., Knorr, A., Hey, R., Ploog, K.H., Radiative coupling of intersubband transitions in GaAs/AlGaAs multiple quantum wells (2006) *Physica e*, 32, pp. 262-265
10. Butscher, S., Knorr, A., Occurrence of intersubband polaronic repellons in a two-dimensional electron gas (2006) *Phys. Rev. L*, 97, p. 197401
11. Yakar, Y., Akir, B., Zmen, A., Calculation of linear and nonlinear optical absorption coefficients of a spherical quantum dot with parabolic potential (2010) *Opt. Commun.*, 283, pp. 1795-1800
12. Zmen, A., Yakar, Y., Akir, B., Atav, ., Computation of the oscillator strength and absorption coefficients for the intersubband transitions of the spherical quantum dot (2009) *Opt. Commun.*, 282, pp. 3999-4004
13. Karabulut, I., Baskoutas, S., Linear and nonlinear optical absorption coefficients and refractive index changes in spherical quantum dots: Effects of impurities, electric field, size, and optical intensity (2008) *J. Appl. Phys.*, 103, p. 073512
14. Bau, N.Q., Trien, H.D., The nonlinear absorption of a strong electromagnetic wave by confined electrons in rectangular quantum wires (2010) *PIERS Proceedings*, pp. 336-341. , Xi'an, China, Mar. 22-26

15. Bau, N.Q., Dinh, L., Phong, T.C., Absorption coefficient of weak electromagnetic waves caused by confined electrons in quantum wires (2007) J. Korean.Phys. Soc., 51, pp. 1325-1330
16. Epstein, E.M., To the theory of nonlinear high frequency conductivity of an electron gas in semiconductors (1970) Sov. Phys. Solid State, 12, pp. 3461-3465
17. Abouelaoualim, D., Electron-confined LO-phonon scattering in GaAs-Al_{0.45}Ga_{0.55}As superlattice (2006) Pramana Journal of Physics, 66, pp. 455-465
18. Gaggero-Sager, M.L., Moreno-Martinez, N., Rodriguez-Vargas, I., Perez-Alvarez, R., Grimalsky, V.V., Mora-Ramos, M.E., Electronic structure as a function of temperature for Si doped quantum wells in GaAs (2007) PIERS Online, 3 (6), pp. 851-854
19. Samuel, E.P., Patil, D.S., Analysis of wavefunction distribution in quantum well biased laser diode using transfer matrix method (2008) Progress in Electromagnetics Research Letters, 1, pp. 119-128
20. Ariza-Flores, A.D., Rodriguez-Vargas, I., Electron subband structure and mobility trends in P-N deltadoped quantum wells in Si (2008) Progress in Electromagnetics Research Letters, 1, pp. 159-165