

Detection of protease inhibitors by a reverse zymography method, performed in a tris(hydroxymethyl)aminomethane-Tricine buffer system

Le Q.T., Katunuma N.

Institute for Health Sciences, Tokushima Bunri University, Yamashiro-cho, Tokushima City, Tokushima 770-8514, Japan; Biotechnology Center, Vietnam National University, Hanoi 144 Xuan Thuy-Cau Giay, Hanoi, Viet Nam

Abstract: A new detecting method for protease inhibitors, especially for low-molecular-weight inhibitors, is reported. Inhibitor samples were separated on a protein substrate-SDS-polyacrylamide gel in a Tris-Tricine buffer system that improves the separation and identification of peptides and low-molecular-weight proteins. After electrophoresis, the gel was incubated with the target proteases to hydrolyze the background protein substrate. The inhibitor bands, which were protected from proteolysis by the target proteases, were stained. Standard low-molecular-weight inhibitors, such as pepstatin A for pepsin or matrix metalloproteases inhibitor I for collagenase, as well as larger inhibitors, such as soybean trypsin inhibitor or aprotinin for trypsin and cystatin C for papain, were demonstrated by this method and showed clear blue inhibitor bands in the white background when the gels were treated with the target proteases. Some significant applications of this method are introduced. This method is an ideal system for discovering new protease inhibitors in small natural samples. ?? 2003 Elsevier Inc. All rights reserved.

Author Keywords: Protease inhibitors; Reverse zymography

Index Keywords: aprotinin; buffer; collagenase; cystatin C; matrix metalloproteinase inhibitor; methane; papain; pepsin A; pepstatin; peptide; proteinase; proteinase inhibitor; soybean trypsin inhibitor; tris(hydroxymethyl)aminomethane tricine; trypsin; unclassified drug; analytic method; article; controlled study; gel; human; hydrolysis; molecular weight; polyacrylamide gel electrophoresis; priority journal; protein analysis; protein degradation; protein determination; separation technique; staining; zymography; Buffers; Electrophoresis, Polyacrylamide Gel; Endopeptidases; Glycine; Protease Inhibitors; Proteins; Staining and Labeling; Tromethamine; Glycine max

Year: 2004

Source title: Analytical Biochemistry

Volume: 324

Issue: 2

Page : 237-240

Cited by: 10

Link: [Scopus Link](#)

Chemicals/CAS: aprotinin, 11004-21-0, 12407-79-3, 50936-63-5, 52229-70-6, 58591-29-0, 9050-74-2, 9075-10-9, 9087-70-1; collagenase, 37288-86-1, 39433-96-0; methane, 74-82-8; papain, 9001-73-4; pepsin A, 9001-75-6; pepstatin, 26305-03-3, 39324-30-6; proteinase inhibitor, 37205-61-1; proteinase, 9001-92-7;

soybean trypsin inhibitor, 9078-38-0; trypsin, 9002-07-7; Buffers; Endopeptidases, EC 3.4.-; Glycine, 56-40-6; Protease Inhibitors; Proteins; tricine, 5704-04-1; Tromethamine, 77-86-1

Correspondence Address: Katunuma, N.; Institute for Health Sciences, Tokushima Bunri University, Yamashiro-cho, Tokushima City, Tokushima 770-8514, Japan; email: katunuma@tokushima.bunri-u.ac.jp

ISSN: 32697

CODEN: ANBCA

DOI: 10.1016/j.ab.2003.09.033

PubMed ID: 14690687

Language of Original Document: English

Abbreviated Source Title: Analytical Biochemistry

Document Type: Article

Source: Scopus

Authors with affiliations:

1. Le, Q.T., Institute for Health Sciences, Tokushima Bunri University, Yamashiro-cho, Tokushima City, Tokushima 770-8514, Japan, Biotechnology Center, Vietnam National University, Hanoi 144 Xuan Thuy-Cau Giay, Hanoi, Viet Nam
2. Katunuma, N., Institute for Health Sciences, Tokushima Bunri University, Yamashiro-cho, Tokushima City, Tokushima 770-8514, Japan

References:

1. Bushell, G.R., Ghosh, P., Hanspal, J.S., Detection of protease inhibitors using substrate-containing sodium dodecyl sulfate-polyacrylamide gel electrophoresis (1983) *Anal. Biochem.*, 132, pp. 288-293
2. Herron, G., Michael, S., Banda, J., Clark, J., Gavrilovic, J., Werb, Z., Secretion of metalloproteinases by stimulated capillary endothelial cells (1986) *J. Biol. Chem.*, 261, pp. 2814-2818
3. Seidl, D.S., Jaffe, W.G., Gonzalez, E., Callejas, A., Microelectrophoretic method for the detection of proteinase inhibitors (1978) *Anal. Biochem.*, 88 (2), pp. 417-424
4. Michcaud, D., Cantin, D., Raworth, A., Vrain, Assessing the stability of cystatin/cystein complex using denaturing gelatin-polyacrylamide gel electrophoresis (1996) *Electrophoresis*, 17, pp. 74-79
5. Schagger, H., Von Jagow, Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of protein in the range from 1 to 100 kDa (1987) *Anal. Biochem.*, 166, pp. 368-379
6. Marcelo, F., Claudia, L., Electrophoretic analysis (Tricine-SDS-PAGE) of bovine caseins (2002) *Acta Farm. Bonaerence*, 21, pp. 57-60
7. Laemmli, U.K., Cleavage of structural protein during the assembly of the head of bacteriophage T4 (1970) *Nature*, 227, pp. 680-685
8. Ohashi, A., Murata, E., Yamamoto, K., Majima, E., Sano, E., Le Q., T., Katunuma, N., New functions of lactoferrin and beta-casein in mammalian milk as cysteine protease inhibitors (2003) *Biochem. Biophys. Res. Commun.*, 306, pp. 98-103
9. Isemura, S., Saitoh, E., Ito, E., Isemura, M., Sanada, K., Cystatin S: A cysteine proteinase inhibitor of human saliva (1984) *J. Biochem. (Tokyo)*, 96, pp. 1311-1314
10. Wojnar, P., Vant Hof, W., Merschak, P., Redl, B., The N-terminal part of recombinant human tear lipocalin/von Ebner's gland protein confers cysteine proteinase inhibition depending on the presence of the entire cystatin-like sequence motifs (2001) *Biol. Chem.*, 382 (10), pp. 1515-1520